

DID YOU KNOW YOU CAN REDUCE NEW HIV & STI CASES IN OREGON?



Oregon experienced a 900% increase in syphilis rates between 2010 and 2021.¹ OHA recommends screening all sexually active individuals under 45 at least once if not tested since January 2021.²



70 - 88% of extragenital chlamydia and gonorrhea cases are missed with urine screening alone in men who have sex with men.³ Resources are available to increase your clinic's capacity to perform extragenital site testing.



Oral HIV Pre-Exposure Prophylaxis (PrEP) options are available for ~ \$30/month.⁴ CDC recommends discussing HIV PrEP with everyone who is sexually active.⁵

Connect with Oregon AETC Clinical Faculty to Take Action

AETC clinical faculty mentor providers to help distill the latest evidence and identify relevant tools and resources to save time and improve patient outcomes. Sessions are brief, no cost, and free of commercial bias.

STI SCREENING & TREATMENT • SYPHILIS & CONGENITAL SYPHILIS •
HIV CLINICAL PREVENTION (PREP & PEP) • HIV DIAGNOSIS & ART INITIATION

Sign Up for a Virtual Mentoring Session Today

Visit oraetc.org/prescriber-support or scan the code to learn more

This opportunity is available to physicians, advanced practitioners, nurses, and pharmacists practicing in the state of Oregon.



¹⁻⁵ Visit tinyurl.com/yc6758kn to view references